

Precision Cosmology and the Density of Baryons in the Universe

Manoj Kaplinghat¹ and Michael S. Turner^{1,2,3}

¹ *Department of Astronomy and Astrophysics
The University of Chicago, 5640 S. Ellis Ave., Chicago, IL 60637, USA*

² *NASA/Fermilab Astrophysics Center
Fermi National Accelerator Laboratory, PO Box 500
Batavia, IL 60510-0500 USA*

³ *Department of Physics, Enrico Fermi Institute
The University of Chicago, Chicago, Illinois 60637 USA*

Big-bang Nucleosynthesis (BBN) and Cosmic Microwave Background (CMB) anisotropy measurements give independent, accurate measurements of the baryon density and can test the framework of the standard cosmology. Early CMB data are consistent with the longstanding conclusion from BBN that baryons constitute a small fraction of matter in the Universe, but may indicate a slightly higher value for the baryon density. We clarify precisely what the two methods determine, and point out that differing values for the baryon density can indicate either an inconsistency or physics beyond the standard models of cosmology and particle physics. We discuss other signatures of the new physics in CMB anisotropy.

Introduction. Just a decade ago the phrase “precision cosmology” would have been an oxymoron. The COBE FIRAS determination of the temperature of the Cosmic Microwave Background (CMB) to four significant figures, $T_0 = 2.725 \pm 0.001$ K [1] (we quote all errors at 1σ) should dispel such thoughts. Cosmologists now foresee a precision era where a flood of high-quality data, from measurements of CMB anisotropy to large-scale structure, pin down cosmological parameters to percent-level precision, decisively testing theories of the early Universe and probing physics at energy scales beyond those accessible in accelerator experiments [2].

Some of the data can also test the consistency of the standard cosmology. In particular, the longstanding big-bang nucleosynthesis determination of the baryon density, will be checked by CMB anisotropy data at one percent or better [3]. The physics underlying the two measurements could hardly be more different: the baryon density at a time of one second determines how complete the conversion of neutrons and protons to tightly-bound ^4He nuclei is, while the baryon density 400,000 years later (at the time of last scattering) determines the amplitude of gravity-driven acoustic oscillations in the baryon-photon fluid.

The determination of the primeval deuterium abundance in a number of high-redshift ($z \sim 2 - 4$) hydrogen clouds [4] coupled with refined predictions of the standard theory of BBN has led to a determination of the baryon density to an accuracy of about 5%, $\Omega_B h^2 = 0.019 \pm 0.00095$ [5]. Very recently, the BOOMERanG [6] and MAXIMA [7] balloon-borne CMB anisotropy experiments have mapped CMB anisotropy with sufficient angular resolution to make the first CMB determinations of the baryon density, $\Omega_B h^2 = 0.032^{+0.005}_{-0.004}$ [8]. While the CMB result has a much larger uncertainty (and depends upon the imposed priors and parameters that are allowed to vary [9]), additional data and new CMB measurements should soon narrow the gap between the two methods.

The early results are encouraging for the consistency of the standard cosmology: the CMB value for the baryon density agrees with the BBN determination within about 2σ and lies far from dynamical determinations of the total mass density, $\Omega_M h^2 = 0.2 \pm 0.04$ (inferred from $\Omega_M = 0.35 \pm 0.07$ [10] and $h = 0.7 \pm 0.07$ [11]). Thus, the CMB measurement strongly supports the case for non-baryonic dark matter, whose linchpin for twenty years has been the discrepancy between the BBN value for Ω_B and dynamical measurements of Ω_M [3,12].

The purpose of our *Letter* is to clarify exactly what is determined, with regard to the baryon density, by BBN and CMB anisotropy and to point out that the two determinations need not agree to be consistent. A discrepancy could very well be the signal of new physics. Here and throughout Ω_B denotes the fraction of critical density in baryons today and $h = H_0/100 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The physical baryon density today, $\rho_B = 1.88 \times 10^{-29} (\Omega_B h^2) \text{ g/cm}^3$.

Comparing apples to apples. In the standard theory of BBN (i.e., isotropic and homogeneous Universe with only the known particle species and the assumption that all three neutrino species are light with mass $\ll 1 \text{ MeV}$, and negligible chemical potentials), the yields of BBN depend only upon the baryon-to-photon ratio η , the neutron mean lifetime and eleven key nuclear cross sections [3,12]. The uncertainties due to the neutron mean lifetime and nuclear data have recently been re-evaluated and significantly reduced [5]. Based upon these predictions and uncertainties, the Burles-Tytler primeval deuterium measurement, $(\text{D}/\text{H})_P = (3.4 \pm 0.3) \times 10^{-5}$, implies a baryon-to-photon ratio $\eta_{\text{BBN}} = (5.1 \pm 0.2) \times 10^{-10}$ [13].

In order to infer the present density of baryons one must convert η_{BBN} to a baryon density at the time of BBN by multiplying by the photon number density, $n_\gamma(\text{BBN}) = 2\zeta(3)T_{\text{BBN}}^3/\pi^2$, and the mean mass per baryon ($\equiv \bar{m}$) and then reduce that density by the vol-

ume increase of the Universe since,

$$\rho_B(\text{today}) = 2\zeta(3)\bar{m}\eta_{\text{BBN}} R_{\text{BBN}}^3 T_{\text{BBN}}^3 / \pi^2 R_0^3, \quad (1)$$

where $R(t)$ is the cosmic scale factor and baryon-number conservation since BBN has been explicitly assumed.

Because we do not know the value of scale factor at the time of BBN *a priori*, we cannot proceed without further assumptions. The standard assumption is adiabaticity: the constancy of the electromagnetic entropy per unit comoving volume, which is proportional to $(RT)^3$ [14], since BBN. Allowing for the possibility that the entropy per unit comoving volume has changed it follows that

$$\rho_B(\text{today}) = \frac{S_{\text{EM}}(\text{BBN})}{S_{\text{EM}}(\text{today})} \eta_{\text{BBN}} \bar{m} n_\gamma(\text{today}). \quad (2)$$

The number density of photons today is related to the present temperature of the CMB, $n_\gamma = (410.5 \pm 0.5) \text{ cm}^{-3}$; dividing by the critical density we arrive at the key equation,

$$\Omega_B h^2 = (0.019 \pm 0.00095) \frac{S_{\text{EM}}(\text{BBN})}{S_{\text{EM}}(\text{today})}. \quad (3)$$

Entropy production after BBN (e.g., by the out-of-equilibrium decay of a massive particle) can increase S_{EM} , which would diminish the BBN prediction for the baryon density. It is also possible to reduce S_{EM} (at the expense of more exotic physics [15]), increasing the BBN prediction for the present baryon density.

The signature of the baryon density in the CMB involves the heights of the “acoustic peaks” in the angular power spectrum. Around the time of last scattering the photon – baryon fluid is undergoing gravity driven acoustic oscillations; Fourier modes caught at maximum compression (odd peaks) or maximum rarefaction (even peaks) produce the highest amplitude temperature fluctuations on the sky, leading to a series of acoustic peaks in the angular power spectrum (see Fig. 1) [16].

The ratio of the heights of the odd and even peaks increases with baryon density; all other cosmological parameters tend to move the heights of the peaks in unison. Thus, determining the baryon density does not suffer from the cosmic degeneracies that affect other parameters, and an ultimate precision of better than one percent can be expected [16].

Converting the baryon density at last scattering (redshift $z_{\text{LS}} \simeq 1100$) to the present involves only a factor of $(1 + z_{\text{LS}})^3$ [17]. While a number of analyses of the BOOMERanG and MAXIMA data have been carried out [9], the current state of affairs is probably fairly represented by the joint analysis of the two teams [8],

$$\Omega_B h^2 = 0.032_{-0.004}^{+0.005}. \quad (4)$$

Changing entropy and its CMB signature. The entropy increase due to the out-of-equilibrium decay (*i.e.*, when

$T \ll m$) of a massive particle relic is given in terms of the particle’s lifetime and mass [14]:

$$r \equiv S_f/S_i \approx 3 m_{\text{Pl}}^{-1/2} Y m \tau^{1/2}, \quad (5)$$

where $Y \equiv n/s$ is the pre-decay abundance (number density per unit comoving entropy density – for a neutrino species, $Y = 0.04$). We have assumed that the decay products thermalize into photons and increase the EM entropy; to avoid distorting the nearly perfect black-body spectrum of the CMB this must occur before about 10^6 sec. In this case the baryon density today is smaller (than $\Omega_B h^2 = 0.019$) by a factor of r .

The increase in entropy has a CMB signature which involves the fact that the energy density in relativistic particles at the time of last scattering is smaller than in the standard scenario. This is because EM entropy production increases the photon-to-neutrino temperature ratio by a factor of $r^{1/3}$ over the standard value, $T_\gamma/T_\nu = r^{1/3}(11/4)^{1/3}$, and thereby decreases the energy in neutrinos at last scattering (which occurs at a fixed photon temperature, $T_{\text{LS}} \simeq 0.3 \text{ eV}$). This decrease in relativistic energy density can be expressed in terms of a (lower) equivalent number of standard neutrino species (see Fig. 2):

$$N_\nu^{\text{equiv}} = 3/r^{4/3}. \quad (6)$$




FIG. 1. The multipole power spectrum of CMB anisotropy for $N_\nu^{\text{equiv}} = 3, 8.3$ (dashed and solid curves respectively) and $\Omega_B h^2 = 0.03$, $\Omega_M h^2 = 0.17$, $\Omega_0 = 1$ and $\Omega_\Lambda = 1 - \Omega_M$. Note the characteristic series of acoustic peaks and the large change in the power spectrum due to the change in N_ν^{equiv} .

More or less energy in relativistic particles at the time of last scattering affects the spectrum of CMB anisotropy by changing the expansion rate and the rate at which the gravitational potentials associated with density perturbations decay. Less relativistic energy increases the sound horizon at last scattering, thereby shifting the acoustic peaks to larger angular scales (smaller ℓ). Less relativistic

energy also depresses the power around the first acoustic peak by diminishing the integrated Sachs-Wolfe effect [16], and decreases the damping of the higher-order acoustic peaks (because the peaks have shifted to larger scales relative to the damping length) as can be seen in Fig. 1.

Two points deserve mention here: (1) smaller N_ν^{equiv} also changes the power spectrum of matter inhomogeneity by shifting the epoch of matter – radiation equality to an earlier time. (2) Only a change in the EM entropy affects the predicted BBN baryon density; decays without EM decay products (*e.g.*, decay to three neutrinos) do not affect the BBN prediction. If the massive-particle decays produce both EM and non-EM entropy, the EM decay products determine r , and the non-EM decay products increase the energy density in relativistic particles and lead to an additional term in Eq. (6).

The value of N_ν^{equiv} will ultimately be determined from CMB anisotropy to a few percent [18]. This provides a cross check on this explanation: if the present baryon density inferred from BBN exceeds that inferred from CMB by a factor of ξ , then N_ν^{equiv} should be $3/\xi^{4/3}$. Of course, the current data is inconsistent with any post-BBN entropy production, and seems to require $r < 1$, *i.e.*, entropy reduction.

Entropy reduction. By invoking more exotic physics it is also possible to reduce the EM entropy. In order to do so without violating the 2nd law of thermodynamics, photons must come into thermal contact with a cooler reservoir (denoted as the X sector) after BBN and before last scattering. Such an idea was previous discussed in a different context [15], and we will briefly discuss its possible relevance here.

The basic idea is simple; owing to the energy dependence of the cross section for X-sector particles to interact with photons (and other familiar particles), the X sector is decoupled (interaction rate per particle Γ less than the expansion rate H) at high temperatures. (This can be achieved provided that $\langle\sigma v\rangle \propto 1/T^n$ with $n > 1$.) At a temperature T_c , between the epoch of BBN and the present, the interactions become rapid enough to quickly establish thermal contact and $T_X = T$, thereby draining entropy from the photons (assuming that the X sector was cooler).

The authors of Ref. [15] have shown that a self-consistent model for the X sector can be constructed, and further, that it is consistent with astrophysical considerations that constrain the interaction of unseen particles with photons (*e.g.*, emission of X sector particles through plasmon processes in red giants and supernovae). With that in mind, let us proceed.

For simplicity, assume that the transfer of energy to X particles proceeds quickly, by thermally populating massless degrees of freedom in the X sector with total statistical weight g_X . The decrease in the EM entropy, $r = S_{\text{EM}}(\text{today})/S_{\text{EM}}(\text{BBN})$, follows from energy conser-

vation:

$$r = [2/(2 + g_X)]^{3/4}. \quad (7)$$

The present *apparent* discrepancy between the BBN and CMB determinations could be explained with $r \simeq 2/3$ ($\Rightarrow g_X \simeq 1.5$).

Provided $T_c \geq T_{LS}$, this scenario also leads to an increase in energy density in relativistic particles at the time of last scattering and a signature in CMB anisotropy as discussed above. The energy increase arises from: (1) the energy density in X-sector particles; and (2) the higher neutrino-to-photon temperature ratio, $T_\nu/T = r^{-1/3}(4/11)^{1/3}$. Again, parameterizing this by the equivalent number of standard neutrino species, it follows that

$$N_\nu^{\text{equiv}} = \frac{4}{7} \left(\frac{4}{11} \right)^{4/3} g_X + 3/r^{4/3}, \quad (8)$$

where the two terms correspond to the two effects just mentioned.

We note that the second term is mandatory and robust – depressing the photon temperature necessarily raises the ratio of the neutrino to photon temperature – and is identical in form to the term that arises in the previous case where the entropy is increased, cf. Eq. (6). The first term is model dependent and would be different if the X sector did not reach the same temperature as the photons or if only the massive degrees of freedom were excited. Finally, for $r = 2/3$, $N_\nu^{\text{equiv}} = 8.3$, which as Fig. 1 shows has a dramatic effect on the spectrum of CMB anisotropy.

Concluding remarks. By means of very different physics fine-scale CMB anisotropy and BBN each have the potential to determine the mean baryon density to percent accuracy or better. Currently, the BBN determination has a precision of 5% and the CMB measurement 15%, and the two disagree at about the 2σ level.

As we have emphasized, a disagreement between the two determinations of the baryon density need not indicate inconsistency: if the BBN baryon density is larger (smaller) by some factor, this could be explained by an increase (decrease) in the EM entropy since the time of BBN by the same factor [19]. If the two baryon densities agree, then one can limit any post BBN electromagnetic entropy change. As discussed, any entropy change also has a distinctive testable signature in CMB anisotropy (see Fig. 2).

We have assumed the standard theory of BBN; relaxing one of its assumptions (*e.g.*, a decaying tau neutrino with a mass of $\mathcal{O}(\text{MeV})$ [20] or large neutrino chemical potentials [21]) can also change the baryon density inferred from the primeval deuterium abundance. Both possibilities have been discussed [22]. These solutions work by speeding up the expansion around the end of BBN through additional energy density in relativistic particles

so that more deuterium remains unburnt. Thus, both solutions also lead to additional relativistic energy at last scattering. It should be noted that the larger expansion rate also increases the helium abundance and hence these solutions require another variable to offset the effect on helium.

A positive chemical-potential for the electron neutrino can achieve this offset. In both the decaying tau neutrino and the large chemical potential scenarios, consistency with all light element abundances can be achieved. We, however, consider the decaying tau neutrino scenario less likely in light of recent results [23] from the SuperKamiokande collaboration which hint at a mass much less than an MeV for ν_τ . To make a comparison between post-BBN entropy change and non-standard BBN we fix our attention on the large chemical-potential scenario. Using ξ to denote the ratio between the actual and the BBN baryon density, we have used the standard BBN code to derive the relation between ξ and the minimum N_ν^{equiv} required, shown in Fig. 2. The ξ - N_ν^{equiv} relation in the non-standard BBN case is distinct from that due to post-BBN entropy decrease.




FIG. 2. The relation between N_ν^{equiv} and ξ ($\equiv 0.019/\Omega_B h^2$) for the non-standard BBN (solid) and for post-BBN entropy change (dashed). Note that for the non-standard BBN case $\xi < 1$ and the plotted value of N_ν^{equiv} is the minimum required; for the post-BBN entropy change the plotted curve is the maximum possible.

For more than two decades big-bang nucleosynthesis has been both a critical test of the standard cosmology and a probe of particle physics (*e.g.*, the limit to the number of light neutrino species) and cosmology (*e.g.*, the baryon density). The cross comparison with the baryon density inferred from CMB anisotropy measurements has opened a new window for testing the cosmological framework and exploring physics beyond the standard model. Should the current discrepancy of about 2σ persist, the resolution may well involve non-standard BBN (neutrino chemical potentials) or new physics (entropy change due to new particles). If so, this would be an even more impressive achievement for cosmology than the BBN limit to the number of neutrino species which itself was ultimately confirmed by laboratory experiment.

ACKNOWLEDGMENTS

We acknowledge support by the DOE at Chicago, and by the DOE and NASA grant NAG 5-7092 at Fermilab.

-
- [1] J.C. Mather *et al.*, *Astrophys. J.* **512**, 511 (1999).
 - [2] See *e.g.*, M.S. Turner & J.A. Tyson, *Rev. Mod. Phys.* **71**, S145 (1999).
 - [3] D.N. Schramm & M.S. Turner, *Rev. Mod. Phys.* **70**, 303 (1998).
 - [4] D. Tytler *et al.*, *Physica Scripta*, **T85**, 12 (2000).
 - [5] S. Burles, K.M. Nollett, J.N. Truran & M.S. Turner, *Phys. Rev. Lett.* **82**, 4176 (1999); K.M. Nollett & S. Burles, *Phys. Rev. D* **61**, 123505 (2000).
 - [6] P. de Bernardis *et al.*, *Nature* **404**, 955 (2000).
 - [7] S. Hanany *et al.*, *Astrophys. J. Lett.*, in press (2000) (astro-ph/0005123).
 - [8] A.H. Jaffe *et al.*, astro-ph/0007333.
 - [9] M. Tegmark & M. Zaldarriaga, *Phys. Rev. Lett.*, **85**, 2240 (2000); A.E. Lange *et al.*, astro-ph/0005004; W. Hu *et al.*, astro-ph/006436; S. Esposito *et al.*, astro-ph/0007419; M. White *et al.*, astro-ph/0004385.
 - [10] See *e.g.*, M.S. Turner, *Physica Scripta* **T85**, 210 (2000), or A. Dekel *et al.*, in *Critical Dialogues in Cosmology*, edited by N. Turok (World Scientific, Singapore, 1997), p. 175.
 - [11] J.R. Mould *et al.*, *Astrophys. J.* **529**, 786 (2000).
 - [12] K.A. Olive, G. Steigman & T.P. Walker, *Phys. Rep.* **333**, 389 (2000).
 - [13] The entropy transferred to photons from $e^\pm$ annihilations decreases the baryon-to-photon ratio by $4/11$. η denotes the post $e^\pm$ -annihilation ratio; the deuterium abundance depends only on this value.
 - [14] E.W. Kolb & M.S. Turner, *The Early Universe*, (Addison-Wesley, Redwood City, CA, 1990), Ch. 5.
 - [15] J.G. Bartlett & L.J. Hall, *Phys. Rev. Lett.* **66**, 541 (1991).
 - [16] W. Hu, N. Sugiyama & J. Silk, *Nature* **386**, 37 (1997).
 - [17] Two implicit assumptions have been made: baryon-number conservation, and the standard theory of recombination which determines z_{LS} precisely. The current data support the standard theory of recombination; future data will test this assumption more precisely.
 - [18] R. Lopez *et al.*, *Phys. Rev. Lett.* **82**, 3952 (1999).
 - [19] It is also possible that post-BBN hadronic and EM showers, which destroy ^4He and produce D, could explain $r < 1$. This does not change N_ν^{equiv} . See *e.g.*, S. Dimopoulos *et al.*, *Astrophys. J.* **330**, 545 (1988).
 - [20] G. Gyuk & M.S. Turner, *Phys. Rev. D* **50**, 6130 (1994); A.D. Dolgov, S.H. Hansen, S. Pastor & D.V. Semikoz, *Nucl. Phys.* **B548**, 385 (1999).
 - [21] H. Kang & G. Steigman, *Nucl. Phys.* **B372**, 494 (1992).
 - [22] J. Lesgourgues & M. Peloso, *Phys. Rev. D* **62**, 81301 (2000); S.H. Hansen & F.L. Villante, *Phys. Lett.* **B486**, 1 (2000); S. Esposito *et al.*, *JHEP* **09**, 038 (2000).

[23] S. Fukuda *et al.*, hep-ex/0009001.